porating a wide range of microlithic tool types including the classic crescentic microlith (Fig. 1a).

REFERENCES

BUTLER, W. H. 1958. Some Previously Unrecorded Aboriginal Artifact Sites Near Perth, Western Australia. W. Aust. Nat., 6 (6).

CAMPBELL, T. D. and NOONE, H. V. V. 1943. Some Aboriginal Campsites in the Woakwine Range Region of the South-East of South Australia. Rec. S. Aust. Mus., 7 (4).

McCARTHY, F. D., BRAMELL, E. and NOONE, H. V. V. 1946. The Stone Implements of Australia. Mem. Aust. Mus., No. 9.

SOME ADDITIONS TO THE ANGIOSPERM FLORA OF LAKESIDE STATION, CUE, WESTERN AUSTRALIA

By KEVIN F. KENNEALLY, Botany Department, University of Western Australia

The appearance of recent taxonomic revisions (Carlquist, 1969; Tindale, 1968; Wilson, 1970) has made it possible to determine to species some of the specimens listed in an annotated list of the flora of Lakeside Station, Cue (Kenneally, 1968). These species are listed below along with the collection numbers cited in the carlier publication.

Mimosaccac

Acacia sp. (Coll. No. 74A) = A, pruinocarpa Tindale.

Eriostemon sp. (Coll. No. 39A) = E. sericeus P. G. Wilson.

Goodeniaceae

Goodenia sp. (Coll. No. 23A) = G. grandiflora Sims var. nicholsonii (F. Mucll.) Krause.

Stylidiaceae

Stylidium sp. (Coll. No. 43A) = S. longibracteatum Carlquist.

Further collections made after the publication of the original list revealed three additional species whilst an examination by Mr. B. R. Maslin of the Western Australian Herbarium of all Acacia material collected at Cue, resulted in the identification of two additional species. These five species are listed below.

Mimosaeeae

Acacia linophylla W. V. Fitzg. Trcc/shrub up to 3 metres, flowers in ovoid heads. On breakaways and plain, common.

A. triptycha F. Muell. ex. Benth. Bushy shrub up to 1 mctre, flowers yellow on simple axillary peduncles. Phyllodes terete, strongly ribbed with curved plumose tip. On breakaways, common.

Euphorbiaccae ·

Pseudantlnus nematophorus F. Mucll. Spreading shrub, branches ending in spines, flowers small and white. On breakaways, not common.

Myoporaccae

Eremophila exilifolia F. Muell. Viscid shrub up to 1 metre, flowers mauve. On breakaways, not common.

E. punicea S. Moore, Shrub up to 1 metrc, flowers intense pink, leaves and calyces densely hairy. On breakaways, not common.

REFERENCES

CARLQUIST, S. 1968. Studies in Stylidiaceae. Aliso, 7: 13-64.

KENNEALLY, K. F. 1968. An annotated list of angiosperms of Lakeside Station, Cue, Western Australia, W. Aust. Nat., 10: 182-189.

TINDALE, M. 1968. Notes on the Australian Taxa of Acacia, No. 1.

Contrib. N.S.W. Nat. Herb., 4: 73-78.

WILSON, P. G. 1970. A taxonomic revision of the genera Crowea, Erios-

temon and Phebalium (Rutaceae). Nuytsia, 1: 1-155.